

Senior Physician, Forensic Psychiatric Service

University of Bern

Falkenplatz 18

CH-3012 Bern

Switzerland

e-mail: Dietrich.vanCalker@fpd.unibe.ch

Retired Guest Researcher

Department of Psychiatry and Psychotherapy, University of Freiburg.

Phone: +49-761-27065820,

fax: +49-761-27069170,

email: dietrich.calker@uniklinik-freiburg.de

Important Publications

Serchov T, Clement HW, Schwarz MK, Iasevoli F, Tosh DK, Idzko M, Jacobson KA, de Bartolomeis A, Normann C, Biber K, van Calker D. Increased Signaling via Adenosine A1 Receptors, Sleep Deprivation, Imipramine, and Ketamine Inhibit Depressive-like Behavior via Induction of Homer1a. *Neuron*. 2015 Aug 5;87(3):549-62

van Calker D, Zobel I, Dykierk P, Deimel CM, Kech S, Lieb K, Berger M, Schramm E. Time course of response to antidepressants: predictive value of early improvement and effect of additional psychotherapy. *J Affect Disord*. 2009 Apr;114(1-3):243-53

Schramm E, van Calker D, Dykierk P, Lieb K, Kech S, Zobel I, Leonhart R, Berger M. An intensive treatment program of interpersonal psychotherapy plus pharmacotherapy for depressed inpatients: acute and long-term results. *Am J Psychiatry*. 2007 May;164(5):768-77.

Willmroth F, Drieling T, Lamla U, Marcushen M, Wark HJ, van Calker D. Sodium-myo-inositol co-transporter (SMIT-1) mRNA is increased in neutrophils of patients with bipolar 1 disorder and down-regulated under treatment with mood stabilizers. *Int J Neuropsychopharmacol*. 2007 Feb;10(1):63-71.

Biber K, Lubrich B, Fiebich BL, Boddeke HW, van Calker D. Interleukin-6 enhances expression of adenosine A(1) receptor mRNA and signaling in cultured rat cortical astrocytes and brain slices. *Neuropsychopharmacology*. 2001 Jan;24(1):86-96.

Lubrich B, van Calker D. Inhibition of the high affinity myo-inositol transport system: a common mechanism of action of antidepressants? *Neuropsychopharmacology*. 1999 Oct;21(4):519-29.

Spleiss O, van Calker D, Schärer L, Adamovic K, Berger M, Gebicke-Haerter PJ. Abnormal G protein alpha(s)- and alpha(i2)-subunit mRNA expression in bipolar affective disorder. Mol Psychiatry. 1998 Nov;3(6):512-20.

Biber, K., Klotz K-N, Berger, M., Gebicke-Härter, P., van Calker, D. Adenosine A1 receptor-mediated activation of phospholipase C in cultured astrocytes depends on the level of receptor expression. J. Neurosci. 1997; 17, 4956-4964

Lubrich B, Patishi Y, Kofman O, Agam G, Berger M, Belmaker RH, van Calker D. Lithium-induced inositol depletion in rat brain after chronic treatment is restricted to the hypothalamus. Mol Psychiatry. 1997 Sep;2(5):407-12

Biber K, Walden J, Gebicke-Härter P, Berger M, van Calker D. Carbamazepine inhibits the potentiation by adenosine analogues of agonist induced inositolphosphate formation in hippocampal astrocyte cultures. Biol Psychiatry. 1996 Oct 1;40(7):563-7.

Greil W, Steber R, van Calker D. The agonist-stimulated accumulation of inositol phosphates is attenuated in neutrophils from male patients under chronic lithium therapy. Biol Psychiatry. 1991 Sep 1;30(5):443-51.

Löffler F, van Calker D, Hamprecht B. Parathyrin and calcitonin stimulate cyclic AMP accumulation in cultured murine brain cells. EMBO J. 1982;1(3):297-302.

van Calker D, Müller M, Hamprecht B. Regulation by secretin, vasoactive intestinal peptide, and somatostatin of cyclic AMP accumulation in cultured brain cells. Proc Natl Acad Sci USA. 1980 Nov;77(11):6907-11.

van Calker D, Müller M, Hamprecht B. Adenosine regulates via two different types of receptors, the accumulation of cyclic AMP in cultured brain cells. J Neurochem. 1979; 33(5):999-1005

van Calker, D., Müller, M. & Hamprecht, B. Adenosine inhibits the accumulation of cyclic AMP in cultured brain cells. Nature 1978; 276, 839-841.

(for a complete list of publications see: van Calker D in PubMed
(<https://www.ncbi.nlm.nih.gov/pubmed>)